

The information in the tables is taken from product datasheets, (checked 31/10/17). As these can change they should always be checked before a product is used. Not all available products are listed. Combination products should only be used if there is a need to treat both worms and fluke.

FLUKE PRODUCTS FOR SHEEP

PRODUCT	TYPE	CONTAINS	MEAT WITHDRAWAL (DAYS)	KILLS			NOTE
				EI	LI	A	
Fasinex 5%	Drench	Triclabendazole	56	✓	✓	✓	EIs from 2 days
Fasinex 100 10%	Drench	Triclabendazole	35	✓	✓	✓	EIs from 2 days
Tribex 5%	Drench	Triclabendazole	56	✓	✓	✓	EIs from 2 days
Triclafas 5%	Drench	Triclabendazole	56	✓	✓	✓	EIs from 2 days
Flukiver	Drench	Closantel	42		✓	✓	Some effect on EIs
Trodax	Injection	Nitroxynil	49		✓	✓	Don't re-treat within 7 weeks
Zanil	Drench	Oxyclozanide	28			✓	Also kills rumen fluke (NL)
COMBINATION FLUKE/WORMERS							
Combinex	Drench	Triclabendazole and Levamisole	56	✓	✓	✓	EIs from 2 days
Cydectin Triclamox	Drench	Triclabendazole and Moxidectin	31	✓	✓	✓	EIs from 2 days
Fasimec Duo	Drench	Triclabendazole and Ivermectin	27	✓	✓	✓	EIs from 2 days
Closamectin	Injection	Closantel and Ivermectin	28		✓	✓	
Supaverem	Drench	Closantel and Mebendazole	65		✓	✓	Some effect on EIs
Mebadown Super	Drench	Closantel and Mebendazole	65		✓	✓	Some effect on EIs
Downland Fluke and Worm	Drench	Oxyclozanide and Levamisole	5			✓	Also kills rumen fluke (NL)
Levafas Diamond	Drench	Oxyclozanide and Levamisole	5			✓	Also kills rumen fluke (NL)
Albex 2.5%	Drench	Albendazole	5			✓	Use at higher dose*
Endospec	Drench	Albendazole	4			✓	Use at higher dose*
Rycoben	Drench	Ricobendazole	3			✓	Use at higher dose*

EI - early immature fluke (around 0- 6 weeks after infection)
LI = late immature fluke (from around 6 to 8 weeks after infection)
A= adult fluke (from around 10-12 weeks after infection)

FLUKE PRODUCTS FOR BEEF CATTLE

PRODUCT	TYPE	CONTAINS	MEAT WITHDRAWAL (DAYS)	KILLS			NOTE
				EI	LI	A	
Fasinex 10%	Drench	Triclabendazole	56	✓	✓	✓	EIs from 2 weeks
Fasinex 100 10%	Drench	Triclabendazole	56	✓	✓	✓	EIs from 2 weeks
Fasinex 240 24%	Drench	Triclabendazole	52	✓	✓	✓	EIs from 2 weeks
Tribex 10%	Drench	Triclabendazole	56	✓	✓	✓	EIs from 2 weeks
Trodax	Injection	Nitroxynil	60		✓	✓	Don't re-treat within 60 days
Flukiver Bovis	Injection	Closantel	77		✓	✓	Don't re-treat within 11 weeks
Zanil	Drench	Oxyclozanide	28			✓	Also kills rumen fluke (NL)
COMBINATION FLUKE/WORMERS							
Combinex	Drench	Triclabendazole and Levamisole	56	✓	✓	✓	EIs from 2 weeks
Cydectin Triclamox	Pour on	Triclabendazole and Moxidectin	143		✓	✓	
Closamectin	Injection	Closantel and Ivermectin	49		✓	✓	
Closamectin	Pour on	Closantel and Ivermectin	28		✓	✓	
Levafas Diamond	Drench	Oxyclozanide and Levamisole	5			✓	Also kills rumen fluke (NL)
Animec Super	Injection	Clorsulon and Ivermectin	66			✓	
Ivomec Super	Injection	Clorsulon and Ivermectin	66			✓	
Virbamec Super	Injection	Clorsulon and Ivermectin	66			✓	
Albex 10%	Drench	Albendazole	14			✓	Use at higher dose
Endospec 10%	Drench	Albendazole	14			✓	Use at higher dose

NL = not specifically licensed for rumen fluke.

EI - early immature fluke (around 0- 6 weeks after infection)
LI = late immature fluke (from around 6 to 8 weeks after infection)
A= adult fluke (from around 10-12 weeks after infection)