

The information in the tables is taken from product datasheets, (checked 22/11/16). As these can change they should always be checked before a product is used. Not all available products are listed. Combination products should only be used if there is a need to treat both worms and fluke.

### FLUKE PRODUCTS FOR SHEEP

| PRODUCT                   | TYPE      | CONTAINS                       | MEAT WITHDRAWAL (DAYS) | KILLS |    |   | NOTE                          |
|---------------------------|-----------|--------------------------------|------------------------|-------|----|---|-------------------------------|
|                           |           |                                |                        | EI    | LI | A |                               |
| Fasinex 5%                | Drench    | Triclabendazole                | 56                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Fasinex 100 10%           | Drench    | Triclabendazole                | 35                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Tribex 5%                 | Drench    | Triclabendazole                | 56                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Triclafas 5%              | Drench    | Triclabendazole                | 56                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Flukiver                  | Drench    | Closantel                      | 42                     |       | ✓  | ✓ | Some effect on EIs            |
| Trodax                    | Injection | Nitroxynil                     | 49                     |       | ✓  | ✓ | Don't re-treat within 7 weeks |
| Zanil                     | Drench    | Oxyclozanide                   | 28                     |       |    | ✓ | Also kills rumen fluke (NL)   |
| COMBINATION FLUKE/WORMERS |           |                                |                        |       |    |   |                               |
| Combinex                  | Drench    | Triclabendazole and Levamisole | 56                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Cydectin Triclamox        | Drench    | Triclabendazole and Moxidectin | 31                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Fasimec Duo               | Drench    | Triclabendazole and Ivermectin | 27                     | ✓     | ✓  | ✓ | EIs from 2 days               |
| Closamectin               | Injection | Closantel and Ivermectin       | 28                     |       | ✓  | ✓ |                               |
| Supaverm                  | Drench    | Closantel and Mebendazole      | 65                     |       | ✓  | ✓ | Some effect on EIs            |
| Mebadown Super            | Drench    | Closantel and Mebendazole      | 65                     |       | ✓  | ✓ | Some effect on EIs            |
| Downland Fluke and Worm   | Drench    | Oxyclozanide and Levamisole    | 5                      |       |    | ✓ | Also kills rumen fluke (NL)   |
| Levafas Diamond           | Drench    | Oxyclozanide and Levamisole    | 5                      |       |    | ✓ | Also kills rumen fluke (NL)   |
| Albex 2.5%                | Drench    | Albendazole                    | 5                      |       |    | ✓ | Use at higher dose*           |
| Allverm 4%                | Drench    | Ricobendazole                  | 3                      |       |    | ✓ | Use at higher dose*           |
| Endospec                  | Drench    | Albendazole                    | 4                      |       |    | ✓ | Use at higher dose*           |
| Rycoben                   | Drench    | Ricobendazole                  | 3                      |       |    | ✓ | Use at higher dose*           |
| Valbazen                  | Drench    | Albendazole                    | 8                      |       |    | ✓ | Use at higher dose*           |

\* Do not use higher dose during tupping or for 1 month after tups are removed.  
NL = not specifically licensed for rumen fluke.

**EI** – early immature fluke (around 0- 6 weeks after infection)  
**LI** = late immature fluke (from around 6 to 8 weeks after infection)  
**A**= adult fluke (from around 10-12 weeks after infection)

**FLUKE PRODUCTS FOR BEEF CATTLE**

| PRODUCT                          | TYPE      | CONTAINS                       | MEAT WITHDRAWAL (DAYS) | KILLS |    |   | NOTE                           |
|----------------------------------|-----------|--------------------------------|------------------------|-------|----|---|--------------------------------|
|                                  |           |                                |                        | EI    | LI | A |                                |
| Fasinex 10%                      | Drench    | Triclabendazole                | 56                     | ✓     | ✓  | ✓ | EIs from 2 weeks               |
| Fasinex 100 10%                  | Drench    | Triclabendazole                | 56                     | ✓     | ✓  | ✓ | EIs from 2 weeks               |
| Fasinex 240 24%                  | Drench    | Triclabendazole                | 52                     | ✓     | ✓  | ✓ | EIs from 2 weeks               |
| Tribex 10%                       | Drench    | Triclabendazole                | 56                     | ✓     | ✓  | ✓ | EIs from 2 weeks               |
| Trodax                           | Injection | Nitroxynil                     | 60                     |       | ✓  | ✓ | Don't re-treat within 60 days  |
| Flukiver Bovis                   | Injection | Closantel                      | 77                     |       | ✓  | ✓ | Don't re-treat within 11 weeks |
| Zanil                            | Drench    | Oxyclozanide                   | 28                     |       |    | ✓ | Also kills rumen fluke (NL)    |
| <b>COMBINATION FLUKE/WORMERS</b> |           |                                |                        |       |    |   |                                |
| Combinex                         | Drench    | Triclabendazole and Levamisole | 56                     | ✓     | ✓  | ✓ | EIs from 2 weeks               |
| Cydectin Triclamox               | Pour on   | Triclabendazole and Moxidectin | 143                    |       | ✓  | ✓ |                                |
| Closamectin                      | Injection | Closantel and Ivermectin       | 49                     |       | ✓  | ✓ |                                |
| Closamectin                      | Pour on   | Closantel and Ivermectin       | 28                     |       | ✓  | ✓ |                                |
| Levafas Diamond                  | Drench    | Oxyclozanide and Levamisole    | 5                      |       |    | ✓ | Also kills rumen fluke (NL)    |
| Animec Super                     | Injection | Clorsulon and Ivermectin       | 66                     |       |    | ✓ |                                |
| Ivomec Super                     | Injection | Clorsulon and Ivermectin       | 66                     |       |    | ✓ |                                |
| Virbamec Super                   | Injection | Clorsulon and Ivermectin       | 66                     |       |    | ✓ |                                |
| Albex 10%                        | Drench    | Albendazole                    | 14                     |       |    | ✓ | Use at higher dose             |
| Endospec 10%                     | Drench    | Albendazole                    | 14                     |       |    | ✓ | Use at higher dose             |

NL = not specifically licensed for rumen fluke.

**EI** – early immature fluke (around 0- 6 weeks after infection)  
**LI** = late immature fluke (from around 6 to 8 weeks after infection)  
**A**= adult fluke (from around 10-12 weeks after infection)